

Null controllability of Kolmogorov-type equations

K. Beauchard

Received: 5 July 2012 / Accepted: 27 March 2013 / Published online: 7 April 2013
© Springer-Verlag London 2013

Abstract We study the null controllability of Kolmogorov-type equations $\partial_t f + v^\gamma \partial_x f - \partial_v^2 f = u(t, x, v)1_\omega(x, v)$ in a rectangle Ω , under an additive control supported in an open subset ω of Ω . For $\gamma = 1$, with periodic-type boundary conditions, we prove that null controllability holds in any positive time, with any control support ω . This improves the previous result by Beauchard and Zuazua (*Ann Inst H Poincaré Anal Non Linéaire* 26:1793–1815, 2009), in which the control support was a horizontal strip. With Dirichlet boundary conditions and a horizontal strip as control support, we prove that null controllability holds in any positive time if $\gamma = 1$ or if $\gamma = 2$ and ω contains the segment $\{v = 0\}$, and only in large time if $\gamma = 2$ and ω does not contain the segment $\{v = 0\}$. Our approach, inspired from Benabdallah et al. (*C R Math Acad Sci Paris* 344(6):357–362, 2007), Lebeau and Robbiano (*Commun Partial Differ Equ* 20:335–356, 1995), is based on two key ingredients: the observability of the Fourier components of the solution of the adjoint system, uniformly with respect to the frequency, and the explicit exponential decay rate of these Fourier components.

Keywords Null controllability · Degenerate parabolic equations · Carleman estimates · Hypoelliptic systems

The author was partially supported by the “Agence Nationale de la Recherche” (ANR), Projet Blanc EMAQS number ANR-2011-BS01-017-01.

K. Beauchard (✉)
CMLS, Ecole Polytechnique, 91128 Palaiseau Cedex, France
e-mail: Karine.Beauchard@math.polytechnique.fr

1 Introduction

1.1 Main result

We consider Kolmogorov-type equations

$$\partial_t f + v^\gamma \partial_x f - \partial_v^2 f = u(t, x, v) 1_\omega(x, v), \quad (t, x, v) \in (0, +\infty) \times \Omega, \quad (1)$$

where $\gamma \in \mathbb{N}^*$, $\Omega = \mathbb{T} \times (-1, 1)$, \mathbb{T} is the 1D-torus, ω is an open subset of Ω , 1_ω is the characteristic function of this set and, $u(t, x, v)$ is a source term located on the subdomain ω . It is a linear control system in which the state is f and the control u is supported in the subset ω .

Depending on the value of γ , we use different boundary conditions in variable v : periodic type boundary conditions when $\gamma = 1$

$$\begin{cases} f(t, x - t, -1) = f(t, x + t, +1), & (t, x) \in (0, +\infty) \times \mathbb{T}, \\ \partial_v f(t, x - t, -1) = \partial_v f(t, x + t, 1), & (t, x) \in (0, +\infty) \times \mathbb{T}, \end{cases} \quad (2)$$

or Dirichlet boundary conditions when $\gamma \in \mathbb{N}^*$

$$f(t, x, -1) = f(t, x, +1) = 0, \quad (t, x) \in (0, +\infty) \times \mathbb{T}. \quad (3)$$

We will also use initial data

$$f(0, x, v) = f_0(x, v), \quad (x, v) \in \Omega. \quad (4)$$

Definition 1 (*Null controllability*) Let $T > 0$ and $\gamma \in \mathbb{N}^*$. System (1)–(2) [resp. System (1)–(3)] is *null controllable in time T* if, for every $f_0 \in L^2(\Omega)$, there exists $u \in L^2((0, T) \times \Omega)$ such that the solution of the Cauchy problem (1)–(2)–(4) [resp. (1)–(3)–(4)] satisfies $f(T, \cdot, \cdot) = 0$.

When $\gamma = 1$ and $\omega = \mathbb{T} \times (a, b)$ for some $a, b \in (-1, 1)$, the null controllability of system (1)–(2) is proved in [5]. The goals of this article are

1. to improve the strategy of [5] to conclude with more general control supports, in the case of periodic-type boundary conditions when $\gamma = 1$ [i.e. for system (1)–(2)],
2. to study the case of Dirichlet boundary conditions [i.e. system (1)–(3)],
3. to give an indication about the possible critical parameter γ for the null controllability (possibly $\gamma = 2$), for system (1)–(3), as for Grushin equations in [4].

The main results of this paper are the following ones.

- Theorem 1** 1. *If $\gamma = 1$ and ω is an open subset of Ω , then the system (1)–(2) is null controllable in any time $T > 0$.*
2. *If $\gamma = 1$ and $\omega = \mathbb{T} \times (a, b)$ with $-1 < a < b < 1$, then the system (1)–(3) is null controllable in any time $T > 0$.*

3. If $\gamma = 2$ and $\omega = \mathbb{T} \times (a, b)$ with $0 < a < b < 1$, then there exists $T^* \geq a^2/2$ such that
 - the system (1)–(3) is null controllable in any time $T > T^*$,
 - the system (1)–(3) is not null controllable in time $T < T^*$.
4. If $\gamma = 2$ and $\omega = \mathbb{T} \times (a, b)$ with $-1 < a < 0 < b < 1$, then the system (1)–(3) is null controllable in any time $T > 0$.

Note that in the third statement, the set $\{v = 0\}$ is not contained in the control location ω , contrary to the fourth case. Theorem 1 emphasizes several behaviors:

1. A sensitivity to boundary conditions (see the asymptotic behavior of Fourier components in Propositions 2, 10 and 17),
2. A finite speed of propagation from $v = a > 0$ to $v = 0$ with $\gamma = 2$ and Dirichlet boundary conditions.

By duality, Theorem 1 is equivalent to observability results for the adjoint system

$$\partial_t g - v^\gamma \partial_x g - \partial_v^2 g = 0, \quad (t, x, v) \in (0, +\infty) \times \Omega, \tag{5}$$

associated to the following boundary conditions when $\gamma = 1$

$$\begin{cases} g(t, x - T + t, -1) = g(t, x + T - t, 1), & (t, x) \in (0, +\infty) \times \mathbb{T}, \\ \partial_v g(t, x - T + t, -1) = \partial_v g(t, x + T - t, 1), & (t, x) \in (0, +\infty) \times \mathbb{T}, \end{cases} \tag{6}$$

or the following ones for $\gamma \in \mathbb{N}^*$

$$g(t, x, -1) = g(t, x, 1) = 0, \quad (t, x) \in (0, +\infty) \times \mathbb{T}. \tag{7}$$

We will also use initial data

$$g(0, x, v) = g_0(x, v), \quad (x, v) \in \Omega. \tag{8}$$

Definition 2 (Observability) Let $T > 0$ and $\gamma \in \mathbb{N}^*$. System (5)–(6) [resp. System (5)–(7)] is *observable in ω in time T* if there exists $C > 0$ such that, for every $g_0 \in L^2(\Omega)$, the solution of the Cauchy problem (5)–(6)–(8) [resp. (5)–(7)–(8)] satisfies

$$\int_{\Omega} |g(T, x, v)|^2 dx dv \leq C \int_0^T \int_{\omega} |g(t, x, v)|^2 dx dv dt.$$

Theorem 2 1. If $\gamma = 1$ and ω is an open subset of Ω , then the system (5)–(6) is observable in ω in any time $T > 0$.

2. If $\gamma = 1$ and $\omega = \mathbb{T} \times (a, b)$ with $0 < a < b < 1$, then the system (5)–(7) is observable in ω in any time $T > 0$.
3. If $\gamma = 2$ and $\omega = \mathbb{T} \times (a, b)$ with $0 < a < b < 1$, then there exists $T^* \geq a^2/2$ such that
 - the system (5)–(7) is observable in ω in any time $T > T^*$,

- the system (5)–(7) is not observable in ω in time $T < T^*$.
4. If $\gamma = 2$ and $\omega = \mathbb{T} \times (a, b)$ with $-1 < a < 0 < b < 1$, then the system (5)–(7) is observable in ω in any time $T > 0$.

Remark 1 Let us emphasize that, when $\gamma = 2$, $\omega = \mathbb{T} \times (a, b)$ with $0 < a < b < 1$ and $T \leq T^*$, then unique continuation holds for system (5)–(7), i.e. any solution g of (5)–(7) satisfies

$$g \equiv 0 \text{ on } (0, T) \times \omega \Rightarrow g \equiv 0 \text{ on } (0, T) \times \Omega$$

(see Proposition 9 for a proof).

1.2 Motivation and bibliographical comments

1.2.1 Null controllability of the heat equation

The null and approximate controllability of the heat equation are essentially well-understood subjects for both linear and semilinear equations, for bounded or unbounded domains (see, for instance, [16, 19, 21–23, 26, 30, 31, 33, 36, 37, 42, 43]) and also with discontinuous (see, e.g. [6, 7, 17, 39]) or singular ([40] and [18]) coefficients.

In particular, the heat equation on a smooth bounded domain Ω of \mathbb{R}^d ($d \in \mathbb{N}^*$) with a source term located on an open subset ω of Ω is null controllable in arbitrarily small time T and with an arbitrarily small control support ω . This result is due, for the case $d = 1$, to Fattorini and Russell [20, Theorem 3.3], and, for $d \geq 2$, to Imanuvilov [28, 29] (see also the book by Fursikov and Imanuvilov [25]) and Lebeau and Robbiano [31]. It is then natural to wonder whether the same result holds for degenerate parabolic equations.

1.2.2 Boundary-degenerate parabolic equations

The null controllability of parabolic equations degenerating on the boundary of the domain in one space dimension is well-understood, much less so in higher dimension. Given $0 < a < b < 1$ and $\gamma > 0$, let us consider the 1D equation

$$\partial_t w + \partial_x(x^{2\gamma} \partial_x w) = u(t, x)1_{(a,b)}(x), \quad (t, x) \in (0, \infty) \times (0, 1),$$

with suitable boundary conditions. Then, null controllability holds if and only if $\gamma \in (0, 1)$ (see [13, 14]), while for $\gamma \geq 1$, the best result one can show is “regional null controllability” (see [12]), which consists in controlling the solution within the domain of influence of the control. Several extensions of the above results are available in one space dimension, see [1, 34] for equations in divergence form [10, 11], for nondivergence form operators, and [9, 24] for cascade systems. Fewer results are available for multidimensional problems, mainly in the case of two dimensional parabolic operators which simply degenerate in the normal direction to the boundary of the space domain, see [15].

1.2.3 Parabolic equations degenerating inside the domain

In [35], the authors study linearized Crocco type equations

$$\begin{cases} \partial_t f + \partial_x f - \partial_{vv} f = u(t, x, v)1_\omega(x, v), & (t, x, v) \in (0, T) \times \mathbb{T} \times (0, 1), \\ f(t, x, 0) = f(t, x, 1) = 0, & (t, x) \in (0, T) \times \mathbb{T}. \end{cases}$$

For a given strict open subset ω of $\mathbb{T} \times (0, 1)$, they prove that null controllability does not hold: the optimal result is regional null controllability. Note that for Kolmogorov Eq. (1), the coupling between the diffusion (in v) and the transport (in x at speed v) generates diffusion both in variables x and v (see Propositions 2, 10 and 17).

In [4], we study Grushin-type equations

$$\begin{cases} \partial_t f - \partial_x^2 f - |x|^{2\gamma} \partial_y^2 f = u(t, x, y)1_\omega(x, y), & (t, x, y) \in (0, T) \times \Omega, \\ f(t, x, y) = 0, & (t, x, y) \in (0, T) \times \partial\Omega, \end{cases} \tag{9}$$

where $\Omega := (-1, 1) \times (0, 1)$, $\omega \subset (0, 1) \times (0, 1)$, and $\gamma > 0$. Here, the parabolic operator degenerates along the line $\{0\} \times (0, 1)$. We prove that

- null controllability holds in any time $T > 0$ when $\gamma \in (0, 1)$,
- null controllability does not hold (whatever $T > 0$) when $\gamma > 1$,
- when $\gamma = 1$ and $\omega = (a, b) \times (0, 1)$ with $0 < a < b < 1$, there exists $T_{min} \geq a^2/2$ such that null controllability holds when $T > T_{min}$ and does not hold when $T < T_{min}$.

Note that contrary to Grushin-type Eq. (9), in Kolmogorov Eq. (1), the parabolic operator degenerates everywhere on the domain.

1.2.4 Null controllability and hypoellipticity

It could be interesting to analyze the connections between null controllability and hypoellipticity.

We recall that a linear differential operator P with C^∞ coefficients in an open set $\Omega \subset \mathbb{R}^d$ is called hypoelliptic if, for every distribution u in Ω , u must be a C^∞ function in every open set where so is Pu . The following sufficient condition (which is also essentially necessary) for hypoellipticity is due to Hörmander (see [27]).

Theorem 3 *Let P be a second order differential operator of the form $P = \sum_{j=1}^r X_j^2 + X_0 + c$, where X_0, \dots, X_r denote first order homogeneous differential operators in an open set $\Omega \subset \mathbb{R}^n$ with C^∞ coefficients and $c \in C^\infty(\Omega)$. Assume that there exists n operators among*

$$X_{j_1}, [X_{j_1}, X_{j_2}], [X_{j_1}, [X_{j_2}, X_{j_3}]], \dots, [X_{j_1}, [X_{j_2}, [X_{j_3}, [\dots, X_{j_k}] \dots]]],$$

where $j_i \in \{0, 1, \dots, r\}$, which are linearly independent at any given point in Ω . Then, P is hypoelliptic.

The Kolmogorov operator $K := v^\gamma \partial_x + \partial_v^2$ satisfies Hörmander condition for every $\gamma \in \mathbb{N}^*$. Indeed, $K = X_0 + X_1^2$ where

$$X_0(x, v) := \begin{pmatrix} v^\gamma \\ 0 \end{pmatrix}, \quad X_1(x, v) := \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

and

$$[X_0, X_1](x, v) = \begin{pmatrix} \gamma v^{\gamma-1} \\ 0 \end{pmatrix}, \quad [X_1, [X_1, X_2]](x, v) = \begin{pmatrix} \gamma(\gamma-1)v^{\gamma-2} \\ 0 \end{pmatrix}.$$

Thus, when $\gamma = 1$, the first iterated Lie bracket is sufficient; whereas when $\gamma = 2$, the second one is the required (at $v = 0$) to satisfy Hörmander's condition.

First, we emphasize that hypoellipticity is not sufficient for unique continuation. For instance, Alinhac and Zuily [2] built a zero order C^∞ -perturbation of the Kolmogorov operator K for which unique continuation does not hold: there exists C^∞ -functions $u(t, x, v)$ and $a(t, x, v)$ on a neighborhood V of 0 in \mathbb{R}^3 such that $Ku + au = 0$, $u(t, x, v) = v(t, x, v) = 0$ when $v < 0$ and $0 \in \text{Supp}(u)$. Therefore, hypoellipticity cannot be sufficient neither for null controllability.

Let us recall that the Grushin operator $G := \partial_x^2 + |x|^{2\gamma} \partial_y^2$ is the hypoelliptic operator of type II associated to the vector field (X_0, X_1) (i.e. $G = X_0^2 + X_1^2$), whereas Kolmogorov operator K is the one of type I (i.e. $K = X_0 + X_1^2$). Both are prototypes of hypoelliptic operators.

For Grushin-type equations, null controllability (with control on a vertical strip) holds only when the first iterated Lie-bracket is sufficient to satisfy Hörmander's condition ($\gamma \in (0, 1]$). For Kolmogorov-type equations, null controllability (with control on a horizontal strip) holds when the two first iterated Lie-brackets are sufficient ($\gamma \in \{1, 2\}$). A general result which relates null controllability of hypoelliptic operators (depending on their type) to the number of iterated Lie brackets that are necessary to satisfy Hörmander's condition would be very interesting, but remains—for the time being—a challenging open problem.

This article also underlines an important influence of the boundary conditions on the validity of null controllability, through the exponential decay rate of Fourier components (see Propositions 2 and 17).

1.3 Structure of the article

In Sect. 2, we state a global Carleman estimate, for 1D heat equations with parameters, which is a preliminary result for the whole article. In Sect. 3, we prove Theorem 1 for $\gamma = 1$ with periodic type boundary conditions. In Sect. 4, we study the well posedness and the Fourier decomposition of the solutions of (1)–(3) when $\gamma \in \{1, 2\}$. In Sect. 5 (resp. 6), we prove Theorem 1 for $\gamma = 2$ (resp. $\gamma = 1$), with Dirichlet boundary conditions.

2 Preliminary

The goal of this section is the statement of a global Carleman estimate for the Fourier components (in x) of the solution of the adjoint system (5). For $n \in \mathbb{Z}$ and $\gamma \in \mathbb{N}^*$, we introduce the operator

$$\mathcal{P}_{n,\gamma}g := \partial_t g + inv^\gamma g - \partial_v^2 g.$$

Proposition 1 *We assume $\gamma \in \mathbb{N}^*$ (resp. $\gamma = 1$). Let a, b be such that $-1 < a < b < 1$. There exist a weight function $\beta \in C^1([-1, 1], \mathbb{R}_+^*)$, positive constants C_1, C_2 such that, for every $n \in \mathbb{Z}, \gamma \in \{1, 2\}, T > 0$ and $g \in C^0([0, T], L^2(-1, 1)) \cap L^2(0, T; H_0^1(-1, 1))$ (resp. $g \in C^0([0, T], L^2(-1, 1)) \cap L^2(0, T; H^1(-1, 1))$) such that $g(t, -1) = g(t, 1)e^{i2n(T-t)}$ and $\partial_v g(t, -1) = \partial_v g(t, +1)e^{i2n(T-t)}$ the following inequality holds*

$$\begin{aligned} & C_1 \int_0^T \int_{-1}^1 \left(\frac{M}{t(T-t)} \left| \frac{\partial g}{\partial v}(t, v) \right|^2 + \frac{M^3}{(t(T-t))^3} |g(t, v)|^2 \right) e^{-\frac{M\beta(v)}{t(T-t)}} dv dt \\ & \leq \int_0^T \int_{-1}^1 |\mathcal{P}_{n,\gamma}g|^2 e^{-\frac{M\beta(v)}{t(T-t)}} dv dt + \int_0^T \int_a^b \frac{M^3}{(t(T-t))^3} |g(t, v)|^2 e^{-\frac{M\beta(v)}{t(T-t)}} dv dt \end{aligned} \tag{10}$$

where $M := C_2 \max\{T + T^2; \sqrt{|n|}T^2\}$.

The proof of this estimate is classical (see [25]): our weight β is the usual one [see (36), (37), (38) and (39)]. We only track carefully the behavior with respect to n of the different constants. For sake of completeness, a proof is reproduced in Appendix, in the case of Dirichlet boundary conditions on g . For periodic-type boundary conditions, one may use a periodic weight function β , as in [5].

3 Proof of Theorem 1 with $\gamma = 1$ and periodic-type boundary conditions

In all this section, we take $\gamma = 1$.

3.1 Well posedness, Fourier decomposition and dissipation

We have the following well posedness result, for the Cauchy-problem (1)–(2).

Proposition 2 *Let $T > 0, f_0 \in L^2(\Omega)$ and $u \in L^2((0, T) \times \Omega)$. There exists a unique solution $f \in C^0([0, T], L^2(\Omega))$ of the Cauchy problem (1)–(2)–(4). Moreover, if $u \equiv 0$, the Fourier components*

$$f_n(t, v) := \int_{\mathbb{T}} f(t, x, v) e^{-inx} dx, \quad t \in (0, +\infty), v \in (-1, 1), n \in \mathbb{Z}$$

satisfy

$$\|f_n(t, \cdot)\|_{L^2(-1,1)} \leq \|f_n(0, \cdot)\|_{L^2(-1,1)} e^{-\frac{n^2 t^3}{12}}, \quad \forall t > 0, n \in \mathbb{Z}.$$

Proof of Proposition 2 The function $h(t, x, v) := f(t, x + vt, v)$ solves a linear equation with coefficients depending only on t and periodic boundary conditions. Thus, we have an explicit expression

$$\begin{aligned} h(t, x, v) &= \sum_{p, n \in \mathbb{Z}} \hat{f}(n, p) e^{-(p\pi)^2 t + np\pi t^2 - n^2 \frac{t^3}{3}} e^{i(nx + p\pi v)} \\ &+ \sum_{p, n \in \mathbb{Z}} \left(\int_0^t \hat{w}(\tau, n, p) e^{(p\pi)^2 \tau - np\pi \tau^2 + n^2 \frac{\tau^3}{3}} d\tau \right) e^{-(p\pi)^2 t + np\pi t^2 - n^2 \frac{t^3}{3}} e^{i(nx + p\pi v)} \end{aligned}$$

where

$$\begin{aligned} \hat{f}(n, p) &:= \int_{\mathbb{T}} \int_{-1}^1 f_0(x, v) e^{-i(nx + p\pi v)} dv dx, \\ \hat{w}(\tau, n, p) &:= \int_{\mathbb{T}} \int_{-1}^1 1_\omega(x + vt, v) u(\tau, x + vt, v) e^{-i(nx + p\pi v)} dv dx. \end{aligned}$$

The dissipation result is a consequence of the relation

$$-(p\pi)^2 t + np\pi t^2 - n^2 \frac{t^3}{3} = -t \left(p\pi - \frac{nt}{2} \right)^2 - \frac{n^2 t^3}{12}.$$

We refer to [5] for more details. □

3.2 Null controllability of initial data with a finite number of Fourier modes

The goal of this section is the proof of the following result.

Proposition 3 *There exists $C > 0$ such that for every $T > 0, N \in \mathbb{N}^*$ and $f_0 \in L^2(\Omega)$ of the form $f_0(x, v) = \sum_{|n| \leq N} \hat{f}_{0,n}(v) e^{inx}$ there exists a control $u \in L^2((0, T) \times \Omega)$ such that the solution of (1)–(2)–(4) satisfies $f(T, \cdot, \cdot) = 0$ and*

$$\|u\|_{L^2((0,T) \times \Omega)} \leq T e^{C(1 + \frac{1}{T} + N)} \|f_0\|_{L^2(\Omega)}.$$

By duality, this null controllability result is equivalent to the following observability inequality.

Proposition 4 *There exists $C > 0$ such that for every $T > 0$, $N \in \mathbb{N}^*$ and $g_0 \in L^2(\Omega)$ of the form $g_0(x, v) = \sum_{|n| \leq N} g_{0,n}(v)e^{inx}$, the solution of (5)–(6)–(8) satisfies*

$$\int_{\Omega} |g(T, x, v)|^2 dx dv \leq T^2 e^{C(1+\frac{1}{T}+N)} \int_0^T \int_{\omega} |g(t, x, v)|^2 dx dv.$$

For the proof of Proposition 4, we need the 2 following ingredients. The first one is a classical inequality, proved, for example, in [31] (see also [32]).

Proposition 5 *Let $c, d \in \mathbb{R}$ be such that $c < d$. There exists $C > 0$ such that for every $N \in \mathbb{N}^*$ and $(b_n)_{|n| \leq N} \in \mathbb{R}^{2N+1}$,*

$$\sum_{n=-N}^N |b_n|^2 \leq e^{CN} \int_c^d \left| \sum_{n=-N}^N b_n e^{inx} \right|^2 dx.$$

The second ingredient is an estimate of the observability constant for the Fourier components of g .

Proposition 6 *Let $a, b \in \mathbb{R}$ be such that $-1 < a < b < 1$. There exists $C > 0$ such that for every $T > 0$, $n \in \mathbb{Z}$, $g_{0,n} \in L^2(-1, 1)$, the solution of*

$$\begin{cases} \partial_t g_n + invg_n - \partial_v^2 g_n = 0, & (t, v) \in (0, +\infty) \times (-1, 1), \\ g_n(t, -1) = g_n(t, +1)e^{i2n(T-t)}, & t \in (0, +\infty), \\ \partial_v g_n(t, -1) = \partial_v g_n(t, +1)e^{i2n(T-t)}, & t \in (0, +\infty), \\ g_n(0, v) = g_{0,n}(v), & v \in (-1, 1), \end{cases} \tag{11}$$

satisfies

$$\int_{-1}^1 |g_n(T, v)|^2 dv \leq T^2 e^{C(1+\frac{1}{T}+\sqrt{|n|})} \int_0^T \int_a^b |g_n(t, v)|^2 dv dt.$$

Proof of Proposition 6 For $t \in (T/3, 2T/3)$, we have

$$\frac{4}{T^2} \leq \frac{1}{t(T-t)} \leq \frac{9}{2T^2}$$

and

$$\int_{-1}^1 |g_n(T, v)|^2 dv \leq \int_{-1}^1 |g_n(t, v)|^2 dv.$$

Thanks to Proposition 1, we get

$$C_1 \frac{64M^3}{T^6} e^{-\frac{9M\beta^*}{2T^2}} \frac{T}{3} \int_0^1 |g_n(T, v)|^2 dv \leq C_3 \int_0^T \int_a^b |g_n(t, v)|^2 dv dt$$

where $\beta^* := \max\{\beta(x); x \in [-1, 1]\}$, $\beta_* := \min\{\beta(x); x \in [-1, 1]\}$ and $C_3 := \max\{x^3 e^{-\beta_* x}; x \in [-1, 1]\}$. Using the inequality $M \geq C_2[T + T^2]$, we get

$$\int_0^1 |g_n(T, v)|^2 dv \leq C_4 T^2 e^{c_1 \frac{M}{T^2}} \int_0^T \int_a^b |g_n(t, v)|^2 dv dt \tag{12}$$

for some constants $c_1, C_4 > 0$ (independent of n, T and $g_{0,n}$).

First case: $\sqrt{|n|} < 1 + \frac{1}{T}$. Then, $M = C_2(T + T^2)$, thus

$$\int_{-1}^1 |g_n(T, v)|^2 dv \leq C_4 T^2 e^{c_1 C_2 (1 + \frac{1}{T})} \int_0^T \int_a^b |g_n(t, v)|^2 dv dt.$$

Second case: $\sqrt{|n|} \geq 1 + \frac{1}{T}$. Then, $M = C_2 \sqrt{|n|} T^2$, thus

$$\int_0^1 |g_n(T, v)|^2 dv \leq C_4 T^2 e^{c_1 C_2 \sqrt{|n|}} \int_0^T \int_a^b |g_n(t, v)|^2 dv dt.$$

This gives the conclusion. □

Now, let us prove Proposition 4, thanks to Propositions 5 and 6.

Proof of Proposition 4 Let $a, b, c, d \in \mathbb{R}$ be such that $a < b, c < d$ and $(c, d) \times (a, b) \subset \omega$. Let g_n be the solution of (11) for $n = -N, \dots, N$. Then $g(t, x, v) = \sum_{|n| \leq N} g_n(t, v) e_n(x)$, where $e_n(x) := e^{inx}$. From the orthogonality of the family

$(e_n)_{n \in \mathbb{Z}}$ in $L^2(\mathbb{T})$, Propositions 6 and 5, we deduce

$$\begin{aligned} \int_{\Omega} g(T, x, v)^2 dx dv &= \sum_{|n| \leq N-1} \int_0^1 |g_n(T, v)|^2 dv \\ &\leq T^2 e^{C(1+\frac{1}{T}+\sqrt{N})} \sum_{|n| \leq N} \int_0^T \int_a^b |g_n(t, v)|^2 dv dt \\ &\leq T^2 e^{C(N+\frac{1}{T}+\sqrt{N})} \int_0^T \int_a^b \int_c^d \left| \sum_{|n| \leq N} g_n(t, v) e_n(x) \right|^2 dx dv dt \\ &\leq T^2 e^{C(N+\frac{1}{T})} \int_0^T \int_{\omega} |g(t, x, v)|^2 dx dv dt \end{aligned}$$

where the constant C may change from line to line. □

3.3 Construction of the control function

The goal of this section is the proof of the statement 1 of Theorem 1. The construction of the control is the one of [7] (itself inspired from [31], see also [32]).

For $n \in \mathbb{Z}$, we define $e_n(x) := e^{inx}$ and $H_n := e_n \otimes L^2(0, 1)$, which is a closed subspace of $L^2(\Omega)$. For $j \in \mathbb{N}$, we define $E_j := \bigoplus_{|n| \leq 2^j} H_n$ and Π_{E_j} , the orthogonal projection from $L^2(\Omega)$ to E_j .

Let $T > 0$ and $f_0 \in L^2(\Omega)$ and let us build a control $u \in L^2((0, T) \times \Omega)$ such that the solution of (1)–(3)–(4) satisfies $f(T) = 0$. Let $\rho \in \mathbb{R}$ with

$$0 < \rho < \frac{1}{3}. \tag{13}$$

Let $K = K(\rho) > 0$ be such that $K \sum_{j=1}^{\infty} 2^{-j\rho} = T$. Let $(a_j)_{j \in \mathbb{N}}$ be defined by $a_0 = 0$ and $a_{j+1} = a_j + 2T_j$, where $T_j := K2^{-j\rho}$ for every $j \in \mathbb{N}$. We now define the control function u in the following way. On $[a_j, a_j + T_j]$, we apply a control u such that $\Pi_{E_j} f(a_j + T_j) = 0$ and

$$\|u\|_{L^2((a_j, a_j+T_j) \times \Omega)} \leq \mathcal{C}_j \|f(a_j)\|_{L^2(\Omega)}$$

where $\mathcal{C}_j := T_j e^{C(2^j + \frac{1}{T_j})}$ (see Proposition 3). Then

$$\|f(a_j + T_j)\|_{L^2(\Omega)} \leq (1 + \sqrt{T_j} \mathcal{C}_j) \|f(a_j)\|_{L^2(\Omega)}.$$

On $[a_j + T_j, a_{j+1}]$, we apply no control to take advantage of the dissipation of the solution proved in Proposition 2

$$\|f(a_{j+1})\|_{L^2(\Omega)} \leq e^{-2^{2j} \frac{\tau_j^3}{12}} \|f(a_j + T_j)\|_{L^2(\Omega)}.$$

Thus, we obtain

$$\|f(a_{j+1})\|_{L^2(\Omega)} \leq e^{\sum_{k=1}^{2j} \ln(1 + \sqrt{T_k} C_k) - 2^{2k} \frac{\tau_k^3}{12}} \|f_0\|_{L^2(\Omega)}.$$

The choice of ρ ensures that the sum in the exponent tends to $-\infty$ when $j \rightarrow +\infty$, this gives $f(T) = 0$. Arguing in the same way, one proves that the control built above belongs to $L^2((0, T) \times \Omega)$.

4 With Dirichlet boundary conditions: well posedness, Fourier decomposition and unique continuation

In this section, $\gamma \in \{1, 2\}$. Let

$$\mathcal{V} := \{f \in C^\infty(\mathbb{T} \times (-1, 1)); \exists K \subset (-1, 1) \text{ compact s.t. } \text{Supp}(f) \subset \mathbb{T} \times K\}.$$

For $f \in \mathcal{V}$, we define

$$|f|_V := \left(\int_{\Omega} |\partial_v f(x, v)|^2 dx dv \right)^{1/2}$$

and $V := \text{Adh}_{|\cdot|_V}(\mathcal{V})$. Observe that $H_0^1(\Omega) \subset V \subset L^2(\Omega)$, thus V is dense in $L^2(\Omega)$. We define the operator A_γ by

$$D(A_\gamma) := \left\{ f \in V; -\partial_v^2 f + v^\gamma \partial_x f \in L^2(\Omega) \right\},$$

$$A_\gamma f := -\partial_v^2 f + v^\gamma \partial_x f.$$

Then $D(A_\gamma)$ is dense in $L^2(\Omega)$, $(A_\gamma, D(A_\gamma))$ is closed and both A_γ and A_γ^* are dissipative, thus $(A_\gamma, D(A_\gamma))$ generates an strongly continuous semigroup $S_\gamma(t)$ of contractions of $L^2(\Omega)$ (see Lumer-Phillips theorem [38, Corollary 4.4, Chapter 1, page 15], or Hille-Yosida theorem [8, Theorem VII.4, page 105]). For every $T > 0$, $u \in L^2((0, T) \times \Omega)$ and $f_0 \in L^2(\Omega)$, the weak solution of (1)–(3)–(4) is

$$f(t) = S_\gamma(t) f_0 + \int_0^t S_\gamma(t-s)(1_\omega u)(s) ds$$

and the following existence and uniqueness result follows.

Proposition 7 *Let $\gamma \in \{1, 2\}$. For every $T > 0$, $u \in L^2((0, T) \times \Omega)$ and $f_0 \in L^2(\Omega)$ there exists a unique weak solution $f \in C^0([0, T], L^2(\Omega)) \cap L^2((0, T), V)$ of (1)–(3)–(4). Moreover, $f(t) \in D(A_\gamma)$ and $\partial_t f(t) \in L^2(\Omega)$ for a.e. $t \in (0, T)$.*

Let us consider a solution of (5)–(7)–(8) in the sense above. Since $g \in C^0([0, T], L^2(\Omega))$, the function $x \mapsto g(t, x, v)$ belongs to $L^2(\mathbb{T})$ for almost every $(t, v) \in [0, +\infty) \times (-1, 1)$, thus, it can be developed in Fourier series of x as follows:

$$g(t, x, v) = \sum_{n \in \mathbb{Z}} g_n(t, v) e^{inx} \quad \text{where} \quad g_n(t, v) := \int_{\mathbb{T}} g(t, x, v) e^{-inx} dx, \quad \forall n \in \mathbb{Z}. \tag{14}$$

Proposition 8 *For every $n \in \mathbb{Z}$, g_n is the unique solution of*

$$\begin{cases} \partial_t g_n - inv^\gamma g_n - \partial_v^2 g_n = 0, & (t, v) \in (0, +\infty) \times (-1, 1), \\ g_n(t, \pm 1) = 0, & t \in (0, +\infty), \\ g_n(0, v) = g_{0,n}(v), & v \in (-1, 1), \end{cases} \tag{15}$$

where $g_{0,n} \in L^2(-1, 1)$ is given by

$$g_{0,n}(v) := \int_{\mathbb{T}} g_0(x, v) e^{-inx} dx, \quad v \in (-1, 1).$$

This result may be proved by following the same steps as in [4, Section 2.2]. Then, the following unique continuation property follows.

Proposition 9 *Let $\gamma \in \{1, 2\}$, $\omega = \mathbb{T} \times (a, b)$ where $0 < a < b < 1$, $T > 0$ and $g \in C^0([0, T], L^2(\Omega)) \cap L^2((0, T), V)$ a solution of (5)–(7). If $g \equiv 0$ on $(0, T) \times \omega$, then $g \equiv 0$ on $(0, T) \times \Omega$.*

Proof of Proposition 9 Let $n \in \mathbb{Z}$ and g_n be defined by (14). Then $g_n \equiv 0$ on $(0, T) \times (a, b)$ and g_n solves (15). Thus, Proposition 1 ensures that $g_n \equiv 0$ on $(0, T) \times (-1, 1)$. Therefore, $g \equiv 0$ on $(0, T) \times \Omega$. □

5 Proof of Theorem 1 when $\gamma = 2$

In all this section, we take $\gamma = 2$ and $\omega = \mathbb{T} \times (a, b)$ where $-1 < a < b < 1$. In the first 4 subsections, we prove the statement 3 of Theorem 1 and in the last subsection, we prove the statement 4.

5.1 Dissipation speed on $(-1, 1)$

The goal of this section is the proof of the following dissipation property.

Proposition 10 *There exists $K, \delta > 0$ such that, for every $n \in \mathbb{Z} - \{0\}$ and $g_{0,n} \in H^1(-1, 1)$, the solution of (15) satisfies*

$$\int_{-1}^1 |g_n(t, v)|^2 dv \leq K e^{-\delta\sqrt{|n|}t} \int_{-1}^1 \left(\frac{1}{\sqrt{|n|}} |\partial_v g_{0,n}(v)|^2 + \sqrt{|n|} |v g_{0,n}(v)|^2 \right) dv, \quad \forall t > 0.$$

The proof of Proposition 10 relies on the following result.

Proposition 11 *There exists $A, B, C, \delta > 0$ with $B^2 < AC$ such that for every $L > 0$ and $h_0 \in H^1(-L, L)$, the solution of*

$$\begin{cases} \partial_\tau h = \partial_y^2 h + iy^2 h, & (\tau, y) \in (0, +\infty) \times (-L, L), \\ h(\tau, \pm L) = 0, & \tau \in (0, +\infty), \\ h(0, y) = h_0(y), & y \in (-L, L), \end{cases} \tag{16}$$

satisfies

$$\mathcal{L}(t) \leq \mathcal{L}(0)e^{-\delta t}, \quad \forall t > 0, \tag{17}$$

where

$$\mathcal{L}(\tau) = \int_{-L}^L \left(|h(\tau, y)|^2 + A |\partial_y h(\tau, y)|^2 - 2B \Im \left[\overline{y h(\tau, y)} \partial_y h(\tau, y) \right] + C |y h(\tau, y)|^2 \right) dy.$$

Proof of Proposition 11 This proof is inspired from [41]. Let $A, B, C > 0$ be such that

$$B^2 < AC \quad \text{and} \quad A^2 + C^2 < \frac{B}{2} \tag{18}$$

(for instance $A = \epsilon \tilde{A}, B = \epsilon \tilde{B}$ and $C = \epsilon \tilde{C}$ for any $\tilde{A}, \tilde{B}, \tilde{C}, \epsilon > 0$ such that $\tilde{B}^2 < \tilde{A}\tilde{C}$ and $\epsilon(\tilde{A}^2 + \tilde{C}^2) < \tilde{B}/2$). Easy computations give

$$\begin{aligned} \frac{1}{2} \frac{d\mathcal{L}}{d\tau} &= -3B \|yh\|^2 - \|\partial_y h\|^2 - C \|y\partial_y h\|^2 - A \|\partial_y^2 h\|^2 \\ &\quad + C \|h\|^2 - 2A \Im \left[\int_{-L}^L y \partial_y \bar{h} h \right] - 2B \Im \left[\int_{-L}^L y \partial_y^2 \bar{h} \partial_y h \right]. \end{aligned}$$

where $\|\cdot\|$ is the usual $L^2((-L, L), \mathbb{C})$ -norm, i.e.

$$\|f\| := \int_{-L}^L |f(y)|^2 dy.$$

Thanks to the following inequalities

$$\begin{aligned}
 C\|h\|^2 &\leq 2C\|yh\|\|\partial_y h\| \leq \frac{B}{2}\|yh\|^2 + \frac{2C^2}{B}\|\partial_y h\|^2, \\
 -2A\Im \left[\int_0^L y\partial_y \bar{h}h \right] &\leq \frac{B}{2}\|yh\|^2 + \frac{2A^2}{B}\|\partial_y h\|^2, \\
 -2B\Im \left[\int_0^L y\partial_y^2 \bar{h}\partial_y h \right] &\leq A\|\partial_y^2 h\|^2 + \frac{B^2}{A}\|y\partial_y h\|^2,
 \end{aligned}$$

we get

$$\frac{1}{2} \frac{d\mathcal{L}}{d\tau} \leq -2B\|yh\|^2 - \left(1 - \frac{2(A^2 + C^2)}{B} \right) \|\partial_y h\|^2.$$

Thanks to (18), there exists $\delta > 0$ (independent of L) such that $\frac{d\mathcal{L}}{d\tau} \leq -\delta\mathcal{L}$, which gives the conclusion. □

Proof of Proposition 10 One may assume that $n > 0$, otherwise, consider \bar{g}_n . In order to simplify the notations, we write g instead of g_n . The function $h(\tau, y)$ defined by

$$g(t, v) = h(\sqrt{nt}, \sqrt[4]{nv})$$

satisfies (16) with $L = \sqrt[4]{n}$ and $h_0(y) := g_{0,n}(y/\sqrt[4]{n})$. From the previous proposition, we know that

$$\begin{aligned}
 \tilde{\mathcal{L}}(t) = \int_{-1}^1 &\left(|g(t, v)|^2 + \frac{A}{\sqrt{n}} |\partial_v g(t, v)|^2 - 2B\Im \left[\overline{vg(t, v)} \partial_v g(t, v) \right] \right. \\
 &\left. + C\sqrt{n} |vg(t, v)|^2 \right) dv
 \end{aligned}$$

satisfies $\tilde{\mathcal{L}}(t) \leq \tilde{\mathcal{L}}(0)e^{-\delta\sqrt{nt}}$. Moreover, using (18) and

$$\|g\|^2 \leq 2\|vg\|\|\partial_v g\| \leq \sqrt{n}\|vg\|^2 + \frac{1}{\sqrt{n}}\|\partial_v g\|^2$$

we get

$$\tilde{\mathcal{L}}(0) \leq \int_{-1}^1 \left(\frac{2A+1}{\sqrt{n}} |\partial_v g_0(v)|^2 + (2C+1)\sqrt{n} |vg_0(v)|^2 \right) dv.$$

Thus,

$$\begin{aligned} \int_{-1}^1 |g_n(t, v)|^2 dv &\leq \tilde{\mathcal{L}}(t) \\ &\leq \tilde{\mathcal{L}}(0)e^{-\delta\sqrt{n}t} \\ &\leq K \int_{-1}^1 \left(\frac{1}{\sqrt{n}} |\partial_v g_0(v)|^2 + \sqrt{n} |v g_0(v)|^2 \right) dv e^{-\delta\sqrt{n}t} \end{aligned}$$

where $K := \max\{2A + 1; 2C + 1\}$. □

5.2 Null controllability in large time T when $0 < a < b < 1$

In this section, we assume $0 < a < b < 1$. Our goal is to prove the existence of a time $T_1 > 0$ such that for every $T > T_1$, the system (5)–(7) is observable in ω in time T . The following uniform observability result gives the conclusion.

Proposition 12 *There exists $T_1, C > 0$ such that for every $T > T_1, n \in \mathbb{Z}$ and $g_{0,n} \in L^2(-1, 1)$, the solution of (15) satisfies*

$$\int_{-1}^1 g_n(T, v)^2 dv \leq C \int_0^T \int_a^b g_n(t, v)^2 dv dt.$$

Proof of Proposition 12 Working as in the proof of Proposition 6, we get

$$\mathcal{C}_3 e^{-c^* \sqrt{n}} \int_{T/3}^{2T/3} \int_{-1}^1 \left(\sqrt{n} |\partial_v g|^2 + n^{3/2} |g|^2 \right) dv dt \leq \mathcal{C}_4 \int_0^T \int_a^b |g|^2 dv dt \tag{19}$$

for n large enough, where $\mathcal{C}_3 := \mathcal{C}_2 \max\{4\mathcal{C}_1; (4\mathcal{C}_1)^3\}$, $c^* := \frac{9}{2}\mathcal{C}_2 \max\{\beta(v); v \in [-1, 1]\}$, $\mathcal{C}_4 := \max\{x^3 e^{-\beta_* x}; x \geq 0\}$ and $\beta_* := \min\{\beta(v); v \in (a, b)\}$. Moreover, thanks to Proposition 10, we have for any $t \in (T/3, 2T/3)$,

$$\begin{aligned} &\int_{-1}^1 \left(\sqrt{n} |\partial_v g(t, v)|^2 + n^{3/2} |g(t, v)|^2 \right) dv \\ &\geq n \int_{-1}^1 \left(\frac{1}{\sqrt{n}} |\partial_v g(t, v)|^2 + \sqrt{n} |v g(t, v)|^2 \right) dv \\ &\geq \frac{n}{K} e^{\delta\sqrt{n}\frac{T}{3}} \int_{-1}^1 |g(T, v)|^2 dv. \end{aligned}$$

Thus,

$$\int_{-1}^1 |g(T)|^2 dv \leq \frac{1}{nT} C_5 e^{[c^* - \delta T/3]\sqrt{n}} \int_0^T \int_a^b |g|^2 dv dt$$

where $C_5 := 3KC_4/C_3$. This gives the conclusion with $T_1 := 3c^*/\delta$. □

5.3 No null controllability when $0 < a < b < 1$ and $T \leq a^2/2$

In this section, we assume $0 < a < b < 1$. The goal of this section is to prove that (5)–(7) is not observable in ω in time $T \leq a^2/2$, which is equivalent to the following non-uniform observability result.

Proposition 13 *Let $T \leq a^2/2$. For every $C > 0$, there exists $n \in \mathbb{Z}$, $g_{0,n} \in L^2(-1, 1)$ such that the solution of (15) satisfies*

$$\int_{-1}^1 g_n(T, v)^2 dv > C \int_0^T \int_a^b g_n(t, v)^2 dv dt.$$

Proof of Proposition 13 First step: Approximate solution Let $\epsilon > 0$ be such that $b < 1 - \epsilon$ and $\theta \in C^\infty(\mathbb{R})$ be such that

$$\text{Supp}(\theta) \subset (-1 - \epsilon, -1 + \epsilon) \cup (1 - \epsilon, 1 + \epsilon), \quad \theta(\pm 1) = 1.$$

For $n \in \mathbb{N}^*$, we define

$$\tilde{g}_n(t, v) := \sqrt[n]{n} \left(e^{-\sqrt{in} \frac{v^2}{2}} - e^{-\frac{\sqrt{in}}{2}} \theta(v) \right) e^{-\sqrt{in}t}$$

where $\sqrt{i} := e^{i\frac{\pi}{4}}$. We have

$$\begin{cases} \partial_t \tilde{g}_n + inv^2 \tilde{g}_n - \partial_v^2 \tilde{g}_n = \sqrt[n]{n} e^{-\frac{\sqrt{in}}{2}} \left(\sqrt{in} \theta + \theta'' - inv^2 \theta \right) e^{-\sqrt{in}t}, \\ \tilde{g}_n(t, \pm 1) = 0. \end{cases}$$

Let g_n be the solution of

$$\begin{cases} \partial_t g_n + inv^2 g_n - \partial_v^2 g_n = 0, & (t, v) \in (0, T) \times (0, 1), \\ g_n(t, \pm 1) = 0, & t \in (0, T), \\ g_n(0, v) = \tilde{g}_n(0, v), & v \in (0, 1). \end{cases}$$

We have

$$\frac{1}{2} \frac{d}{dt} \|(\tilde{g}_n - g_n)(t)\|_{L^2(-1,1)}^2 = -\|\partial_v(\tilde{g}_n - g_n)(t)\|_{L^2(-1,1)}^2 + \Re \left(\int_{-1}^1 \sqrt[8]{ne^{-\frac{\sqrt{in}}{2}}} \left(\sqrt{in}\theta(v) + \theta''(v) - inv^2\theta(v) \right) e^{-\sqrt{int}} \overline{(\tilde{g}_n - g_n)(t, v)} dv \right).$$

Thanks to Poincare and Cauchy-Schwarz inequalities, we get

$$\frac{d}{dt} \|(\tilde{g}_n - g_n)(t)\|_{L^2(-1,1)} \leq -C_P \|(\tilde{g}_n - g_n)(t)\|_{L^2(-1,1)} + C_1 n^{9/8} e^{-\frac{\sqrt{2n}}{4}} e^{-\frac{\sqrt{2nt}}{2}}$$

where C_P is the Poincare constant on $(-1, 1)$ and C_1 is a positive constant that depends only on θ . Thus

$$\|(\tilde{g}_n - g_n)(t)\|_{L^2(-1,1)} \leq C_2 n^{5/8} e^{-\frac{\sqrt{2n}}{4}}, \quad \forall t \geq 0, \tag{20}$$

where $C_2 > 0$ does not depend on n .

Second step: Conclusion Let $T \leq a^2/2$. Working by contradiction, we assume that there exists $C_T > 0$ such that for every $n \in \mathbb{N}^*$,

$$\int_{-1}^1 g_n(T, v)^2 dv \leq C_T \int_0^T \int_a^b g_n(t, v)^2 dv dt.$$

Thanks to the triangular inequality and (20), we deduce that

$$\begin{aligned} \|\tilde{g}_n(T)\|_{L^2(-1,1)} &\leq \left(C_T \int_0^T \int_a^b |\tilde{g}_n|^2 dv dt \right)^{1/2} + \|(\tilde{g}_n - g_n)(T)\|_{L^2(-1,1)} \\ &\quad + \left(C_T \int_0^T \int_a^b |\tilde{g}_n - g_n|^2 dv dt \right)^{1/2} \\ &\leq \left(C_T \int_0^T \int_a^b |\tilde{g}_n|^2 dv dt \right)^{1/2} + (1 + \sqrt{TC_T}) C_2 n^{5/8} e^{-\frac{\sqrt{2n}}{4}}. \end{aligned}$$

However, there exists $C_3, C_4 > 0$ such that when $n \rightarrow +\infty$,

$$\begin{aligned} \|\tilde{g}_n(T)\|_{L^2} &\sim C_3 e^{-\frac{\sqrt{2nT}}{2}} \\ \left(\int_0^T \int_a^b |\tilde{g}_n(t, v)|^2 dv dt \right)^{1/2} &\sim \frac{C_4}{n^{3/8}} e^{-\frac{\sqrt{2na^2}}{4}}, \end{aligned}$$

which gives a contradiction. □

5.4 End of the proof of Theorems 1.3 and 2.3

Let us consider $\gamma = 2$ and $\omega = \mathbb{T} \times (a, b)$ with $0 < a < b < 1$. From Proposition 12 and Bessel Parseval equality, we know that system (5)–(7) is observable in ω in any time $T > T_1$. From Proposition 13, we deduce that for any time $T \leq \frac{a^2}{2}$, (5)–(7) is not observable in ω in time T . Thus, the quantity

$$T^* := \inf\{T > 0; \text{ system (5)(7) is observable in } \omega \text{ in time } T\}$$

is well-defined and belongs to $[\frac{a^2}{2}, +\infty)$. Clearly, observability in some time T_{\sharp} implies observability in any time $T > T_{\sharp}$, so

- for every $T > T^*$, (5)–(7) is observable in ω in time T ,
- for every $T < T^*$, (5)–(7) is not observable in ω in time T .

5.5 Null controllability in any time $T > 0$ when $a < 0 < b$

In this section, $\omega = \mathbb{T} \times (-a, a)$ where $a > 0$. We fix $\beta \in (0, a)$. Our goal is the proof of the statement 4 of Theorem 1, thanks to a cut-off argument.

5.5.1 Preliminary

We define $\Omega_1 := \mathbb{T} \times (\beta, 1)$ and $\omega_1 := \mathbb{T} \times (\beta, a)$, and we consider the system

$$\begin{cases} \partial_t f + v^2 \partial_x f - \partial_v^2 f = u(t, x, v) 1_{\omega_1}(x, v), & (t, x, v) \in (0, T) \times \Omega_1, \\ f(t, x, \beta) = f(t, x, 1) = 0, & (t, x) \in (0, T) \times \mathbb{T}, \end{cases} \quad (21)$$

The goal of this section is the proof of the following result.

Proposition 14 *The system (21) is null controllable in any time $T > 0$.*

As in Sect. 5.2, this is equivalent to the following observability result.

Proposition 15 *There exists $C > 0$ such that for every $n \in \mathbb{Z}$ and $g_{0,n} \in L^2(\beta, 1)$, the solution of*

$$\begin{cases} \partial_t g_n - i n v^2 g_n - \partial_v^2 g_n = 0, & (t, v) \in (0, +\infty) \times (\beta, 1), \\ g_n(t, \beta) = g_n(t, 1) = 0, & t \in (0, +\infty), \\ g_n(0, v) = g_{0,n}(v), & v \in (\beta, 1), \end{cases} \quad (22)$$

satisfies

$$\int_{\beta}^1 |g_n(T, v)|^2 dv \leq C \int_0^T \int_{\beta}^a |g_n(t, v)|^2 dv dt.$$

For the proof of Proposition 15, we need the following dissipation result.

Proposition 16 *There exists $K, \delta > 0$ such that for every $n \in \mathbb{Z}$, $g_{0,n} \in H^1(\beta, 1)$, the solution of (22) satisfies*

$$\int_{\beta}^1 |g_n(t, v)|^2 dv \leq K e^{-\delta|n|^{2/3}t} \int_{\beta}^1 \left(|g_{0,n}(v)|^2 + \frac{1}{|n|^{2/3}} |\partial_v g_{0,n}(v)|^2 \right) dv.$$

Proof of Proposition 16 One may assume that $n > 0$, otherwise consider $\overline{g_n}$. To simplify the notations, we write g instead of g_n . Let $B, C > 0$ be such that

$$B^2 < 2C \quad \text{and} \quad 3B > 4C^2 \tag{23}$$

(for instance $B = \sqrt{C}$ with $C > 0$ small enough). The function

$$\mathcal{L}(t) := \int_{\beta}^1 \left(|g(t, v)|^2 + \frac{B}{n^{1/3}} \Im \left[v g(t, v) \overline{\partial_v g(t, v)} \right] + \frac{C}{n^{2/3}} |\partial_v g(t, v)|^2 \right) dv$$

satisfies

$$\begin{aligned} \frac{1}{2} \frac{d\mathcal{L}}{dt} &= -\frac{3Bn^{2/3}}{2} \|vg\|^2 - \|\partial_v g\|^2 - \frac{C}{n^{2/3}} \|\partial_v^2 g\|^2 \\ &\quad - 2Cn^{1/3} \Im \left[\int_{\beta}^1 v g \partial_v \overline{g} \right] - \frac{B}{n^{1/3}} \Im \left[\int_{\beta}^1 v \partial_v g \partial_v^2 \overline{g} \right]. \end{aligned}$$

Thanks to

$$\begin{aligned} -2Cn^{1/3} \Im \left[\int_{\beta}^1 v g \partial_v \overline{g} \right] &\leq \frac{1}{2} \|\partial_v g\|^2 + 2C^2 n^{2/3} \|vg\|^2, \\ -\frac{B}{n^{1/3}} \Im \left[\int_{\beta}^1 v \partial_v g \partial_v^2 \overline{g} \right] &\leq \frac{C}{n^{2/3}} \|\partial_v^2 g\|^2 + \frac{B^2}{4C} \|\partial_v g\|^2, \end{aligned}$$

we get

$$\frac{1}{2} \frac{d\mathcal{L}}{dt} \leq -\left(\frac{3B}{2} - 2C^2 \right) n^{2/3} \beta^2 \|g\|^2 - \left(\frac{1}{2} - \frac{B^2}{4C} \right) \|\partial_v g\|^2.$$

Thanks to (23), there exists $\delta > 0$ (independent of n) such that $\frac{d\mathcal{L}}{dt} \leq -\delta n^{2/3} \mathcal{L}$, which gives the conclusion. □

Proof of Proposition 15 Working as in the proof of Proposition 12, we get

$$C_3 e^{-c^* \sqrt{n}} \int_{T/3}^{2T/3} \int_{\beta}^1 \left(\sqrt{n} |\partial_v g|^2 + n^{3/2} |g|^2 \right) dv dt \leq C_4 \int_0^T \int_{\beta}^a |g|^2 dv dt \tag{24}$$

for n large enough. Moreover, thanks to Proposition 16, we have, for any $t \in (T/3, 2T/3)$,

$$\begin{aligned} & \int_{\beta}^1 \left(\sqrt{n} |\partial_v g(t, v)|^2 + n^{3/2} |g(t, v)|^2 \right) dv \\ & \geq n^{7/6} \int_{\beta}^1 \left(\frac{1}{n^{2/3}} |\partial_v g(t, v)|^2 + |g(t, v)|^2 \right) dv \\ & \geq \frac{n^{7/6}}{K} e^{\delta n^{2/3} \frac{T}{3}} \int_{\beta}^1 |g(T, v)|^2 dv. \end{aligned}$$

Thus,

$$\int_{\beta}^1 |g(T)|^2 dv \leq \frac{C_5}{n^{7/6} T} e^{[c^* \sqrt{n} - \delta n^{3/2} T/3]} \int_0^T \int_{\beta}^a |g|^2 dv dt$$

where $C_5 := 3K C_4 / C_3$, which gives the conclusion. □

5.5.2 Cut-off strategy

Let $\xi_i \in C^\infty(\mathbb{R})$ for $i = 1, 2, 3$ such that $0 \leq \xi_i \leq 1$ and

$$\begin{cases} \xi_1 + \xi_2 + \xi_3 \equiv 1 \\ \xi_1(v) = 0 \text{ if } v \leq \beta, \quad \xi_1(v) = 1 \text{ if } v \geq a \\ \xi_2(v) = 1 \text{ if } v \leq -a, \quad \xi_2(v) = 0 \text{ if } v \geq -\beta \\ \xi_3(v) = 1 \text{ if } |v| \leq \beta, \quad \xi_3(x) = 0 \text{ if } |v| \geq a \end{cases} \tag{25}$$

By Proposition 14, there exists $u_1 \in L^2((0, T) \times \Omega_1)$ such that the solution of

$$\begin{cases} \partial_t f_1 + v^2 \partial_x f_1 - \partial_v^2 f_1 = u_1(t, x, v) 1_{\omega_1}(x, v), & (t, x, v) \in (0, T) \times \Omega_1, \\ f_1(t, x, \beta) = f_1(t, x, 1) = 0, & (t, x) \in (0, T) \times \mathbb{T}, \\ f_1(0, x, v) = f_0(x, v), & (x, v) \in \Omega_1, \end{cases} \tag{26}$$

satisfies $f_1(T, \cdot, \cdot) = 0$. Similarly, let $\Omega_2 := \mathbb{T} \times (-1, -\beta)$ and $\omega_2 := \mathbb{T} \times (-a, -\beta)$; there exists $u_2 \in L^2((0, T) \times \Omega_2)$ such that the solution of

$$\begin{cases} \partial_t f_2 + v^2 \partial_x f_2 - \partial_v^2 f_2 = u_2(t, x, v) 1_{\omega_2}(x, v), & (t, x, v) \in (0, T) \times \Omega_2, \\ f_2(t, x, -1) = f_2(t, x, -\beta) = 0, & (t, x) \in (0, T) \times \mathbb{T}, \\ f_2(0, x, v) = f_0(x, v), & (x, v) \in \Omega_2, \end{cases} \tag{27}$$

satisfies $f_2(T, \cdot, \cdot) = 0$. Finally, let $\Omega_3 := \mathbb{T} \times (-\beta, \beta)$ and f_3 be the solution of

$$\begin{cases} \partial_t f_3 + v^2 \partial_x f_3 - \partial_v^2 f_3 = 0, & (t, x, v) \in (0, T) \times \Omega_3, \\ f_3(t, x, \pm\beta) = 0, & (t, x) \in (0, T) \times \mathbb{T}, \\ f_3(0, x, v) = f_0(x, v), & (x, v) \in \Omega_3. \end{cases} \tag{28}$$

We extend f_j and u_j by zero on $(0, T) \times [\Omega - \Omega_j]$ for $j = 1, 2, 3$. Then, the functions

$$\begin{aligned} f(t, x, v) &:= \xi_1(v) f_1(t, x, v) + \xi_2(v) f_2(t, x, v) + \frac{T-t}{T} \xi_3(v) f_3(t, x, v), \\ u(t, x, v) &:= \sum_{j=1}^2 \xi_j(v) u_j(t, x, v) 1_{\omega_j}(x, v) - \frac{1}{T} \xi_3(v) f_3(t, x, v) \\ &\quad - \sum_{j=1}^3 \left(2\xi_j'(v) \partial_v f_j(t, x, v) + \xi_j''(v) f_j(t, x, v) \right) \end{aligned}$$

solve (1)–(3)–(4), as well as $f(T, \cdot, \cdot) = 0$ on Ω . □

6 Proof of Theorem 1 with $\gamma = 1$ and Dirichlet boundary conditions

In all this section, we take $\gamma = 1$ and $\omega = \mathbb{T} \times (a, b)$ where $-1 < a < b < 1$. Our goal is the proof of the statement 2 of Theorem 1. The strategy is the same as in the previous sections, it relies on the following dissipation property.

Proposition 17 *There exists $K, \delta > 0$ such that for every $n \in \mathbb{Z} - \{0\}$ and $g_{0,n} \in H^1(-1, 1)$, the solution of (15) satisfies*

$$\|g_n(t)\|_{L^2(-1,1)} \leq K e^{-\delta|n|^{2/3}t} \|g_{0,n}\|_{H^1(-1,1)}, \quad \forall t > 0. \tag{29}$$

Moreover, the power “2/3” in the exponential rate is optimal as $n \rightarrow +\infty$, and necessarily $\delta \leq \frac{\mu}{2}$, where $-\mu$ is the first zero (from the right) of Airy function in the half line $(-\infty, 0)$.

The proof of Proposition 17 relies on the following result.

Proposition 18 *There exists $B, C, \delta > 0$ with $B^2 < C$ such that for every $L > 0$ and $h_0 \in L^2(-L, L)$, the solution of*

$$\begin{cases} \partial_\tau h = \partial_y^2 h + iyh, & (\tau, y) \in (0, +\infty) \times (-L, L), \\ h(\tau, \pm L) = 0, & \tau \in (0, +\infty), \\ h(0, y) = h_0(y), & y \in (-L, L), \end{cases} \tag{30}$$

satisfies

$$\mathcal{L}(\tau) \leq \mathcal{L}(0) e^{-\delta\tau}, \quad \forall \tau > 0, \tag{31}$$

where

$$\mathcal{L}(\tau) := \int_{-L}^L \left(|h(\tau, y)|^2 - 2B\Im \left[\overline{h(\tau, y)} \partial_y h(\tau, y) \right] + C |\partial_y h(\tau, y)|^2 \right) dy.$$

Proof of Proposition 18 Let $B, C > 0$ be such that

$$4B^2 < C \quad \text{and} \quad 2C^2 < B \tag{32}$$

(for instance, $B = \sqrt{C}/3$ with $C > 0$ small enough). Easy computations show that

$$\frac{1}{2} \frac{d\mathcal{L}}{d\tau} = -B \|h\|^2 - C\Im \left[\int_{-L}^L h \partial_y \bar{h} \right] - \|\partial_y h\|^2 - 2B\Im \left[\int_{-L}^L \partial_y^2 \bar{h} \partial_y h \right] - C \|\partial_y^2 h\|^2.$$

Thanks to

$$\begin{aligned} -2B\Im \left[\int_{-L}^L \partial_y^2 \bar{h} \partial_y h \right] &\leq C \|\partial_y^2 h\|^2 + \frac{B^2}{C} \|\partial_y h\|^2, \\ -C\Im \left[\int_{-L}^L h \partial_y \bar{h} \right] &\leq \frac{B}{2} \|h\|^2 + \frac{C^2}{2B} \|\partial_y h\|^2 \end{aligned}$$

we get

$$\frac{d\mathcal{L}}{d\tau} \leq -\frac{B}{2} \|h\|^2 - \left(1 - \frac{B^2}{C} - \frac{C^2}{2B} \right) \|\partial_y h\|^2.$$

Thanks to (32), there exists $\delta > 0$ such that $d\mathcal{L}/dt \leq -\delta\mathcal{L}$, which gives the conclusion. □

Proof of Proposition 17 First step Proof of (29): One may assume that $n > 0$, otherwise consider \bar{g}_n . In order to simplify the notation, we write g instead of g_n . The function $h(\tau, y)$ defined by

$$g(t, v) = h \left(n^{\frac{2}{3}} t, n^{\frac{1}{3}} v \right)$$

satisfies (30) with $L = n^{\frac{1}{3}}$ and $h_0(y) := g_{0,n}(y/n^{\frac{1}{3}})$. From the previous proposition, we know that

$$\tilde{\mathcal{L}}(t) := \int_{-1}^1 \left(|g(t, v)|^2 - \frac{2B}{n^{\frac{1}{3}}} \Im \left[\overline{g(t, v)} \partial_v g(t, v) \right] + \frac{C}{n^{\frac{2}{3}}} |\partial_v g(t, v)|^2 \right) dv$$

satisfies $\tilde{\mathcal{L}}(t) \leq \tilde{\mathcal{L}}(0)e^{-\delta n^{\frac{2}{3}}t}$. Moreover, using $B^2 < C$ we get

$$\begin{aligned} \int_{-1}^1 |g_n(t, v)|^2 dv &\leq \left(1 - \frac{B^2}{C}\right)^{-1} \tilde{\mathcal{L}}(t) \\ &\leq \left(1 - \frac{B^2}{C}\right)^{-1} \tilde{\mathcal{L}}(0)e^{-\delta n^{\frac{2}{3}}t} \\ &\leq K e^{-\delta n^{2/3}t} \|g_{0,n}\|_{H^1}^2. \end{aligned}$$

for some constant $K > 0$.

Second step Proof of the optimality. First, let us recall that the function

$$\varphi(y) := \text{Ai}\left(e^{i\frac{\pi}{6}}y + \mu\right),$$

satisfies

$$\begin{cases} -\varphi''(y) + iy\varphi(y) = \lambda\varphi(y), & y \in (0, +\infty) \\ \varphi(0) = 0, \end{cases} \tag{33}$$

$$|\varphi(y)| \leq \frac{C}{y^{1/4}} e^{-\frac{\sqrt{2}}{3}y^{3/2}}, \quad \forall y \in (0, +\infty), \tag{34}$$

where $\lambda := -e^{i\frac{2\pi}{3}}\mu$, $C > 0$ (see [3, formulas (2.12) and (A.12)]).

Working by contradiction, we assume that there exists $T > 0$, $n^* \in \mathbb{N}^*$, $(r_n)_{n \in \mathbb{Z}} \in (0, +\infty)^{\mathbb{Z}}$ such that

- any solution of (15) satisfies

$$\|g_n(t)\|_{L^2(-1,1)} \leq K e^{-r_n t} \|g_{0,n}\|_{H^1(-1,1)}, \quad \forall t \in [0, T], n \in \mathbb{Z},$$

- $r_n > (\lambda_r + \delta)|n|^{2/3}$, $\forall |n| \geq n^*$, where $\lambda_r := \Re(\lambda) = |\mu|/2$ and $\delta > 0$.

Let us consider $n \in \mathbb{N}^*$ and $\theta \in C_c^\infty(\mathbb{R})$ such that $\theta(\pm 1) = 1$. Thanks to (33), the function

$$\tilde{g}_n(t, v) := \left(\varphi\left(n^{\frac{1}{3}}(v+1)\right) - \varphi\left(2n^{\frac{1}{3}}\right)\theta(v)\right) e^{-(\lambda n^{2/3} - in)t}$$

satisfies

$$\begin{cases} \partial_t \tilde{g}_n + inv\tilde{g}_n - \partial_v^2 \tilde{g}_n = F_n, & (t, v) \in (0, +\infty) \times (-1, 1), \\ \tilde{g}_n(t, \pm 1) = 0, & t \in (0, +\infty), \end{cases}$$

where

$$F_n(t, v) := \varphi\left(2n^{\frac{1}{3}}\right) \left(\left[\lambda n^{2/3} - in(v+1)\right]\theta(v) + \theta''(v)\right) e^{-(\lambda n^{2/3} - in)t}.$$

Thanks to (34), there exists $C_1 > 0$ such that for every $n \in \mathbb{N}^*$ and $t \in \mathbb{R}$,

$$\|F_n(t)\|_{H^1} \leq C_1 n^{\frac{11}{12}} e^{-\frac{4\sqrt{n}}{3} - \lambda_r n^{\frac{2}{3}} t}.$$

Thanks to the Duhamel formula and (34), we get the following inequality, for every $n \geq n^*$ and $t \in [0, T]$

$$\|\tilde{g}_n(t)\|_{L^2} \leq K e^{-r_n t} \|\tilde{g}_n(0)\|_{H^1} + C_2 n^{\frac{11}{12}} e^{-\frac{4\sqrt{n}}{3}} \int_0^t e^{-r_n(t-s) - \lambda_r n^{\frac{2}{3}} s} ds,$$

where $C_2 > 0$. Thus, there exists a constant $C_3 > 0$ such that for every $n \geq n^*$ and $t \in [0, T]$

$$e^{-\lambda_r n^{\frac{2}{3}} t} \leq C_3 \left(e^{-r_n t} + R(n) e^{-\frac{4\sqrt{n}}{3}} e^{-\lambda_r n^{\frac{2}{3}} t} \right),$$

where R is a rational fraction. We get a contradiction by considering the limit $n \rightarrow +\infty$ (with a fixed $t \in (0, T)$). This ends the proof of the optimality. \square

Remark 2 The optimality of $n^{2/3}$ in the exponential rate shows that we cannot expect to prove the null controllability of (1)–(3) in the same way as we did for (1)–(2). Indeed, the dissipation in $e^{-\delta n^{2/3} t}$ is not sufficient to compensate for the constant e^{Cn} of Proposition 5. Therefore, with Dirichlet boundary conditions, if the null controllability holds with arbitrarily small control supports ω , the proof requires another strategy.

7 Conclusion and open problems

In this article, we have studied the null controllability of Kolmogorov type Eq. (1), with $\gamma \in \{1, 2\}$, in the rectangle $\Omega = \mathbb{T} \times (-1, 1)$, with a distributed control localized on an open subset ω of Ω .

The following questions are still open.

1. When $\gamma > 2$, does null controllability hold? In [4], the proof of the non-uniform observability relies on a comparison argument (maximum principle), which cannot be used here because the 1D heat equation has complex valued coefficients.
2. When $\gamma = 2$, what is the value of the minimal time T^* ? We conjecture that $T^* = a^2/2$.
3. What happens for $\gamma \in (1, 2)$? (with v replaced by $|v|$)
4. With $\gamma = 1$ and Dirichlet boundary conditions in v , does null controllability hold with an arbitrary control support ω ?
5. Is it possible to extend these results to multidimensional configurations? The technique of this paper should possibly extend to cylindrical domains of the form $\mathbb{T} \times (-1, 1)^m$. However, the generalization to more general configurations or boundary controls, is widely open.

Acknowledgments The author warmly thanks Philippe Gravejat, François Golse and Thierry Paul for stimulating discussions.

Appendix: Proof of Proposition 1

Let a', b' be such that $-1 < a < a' < b' < b < 1$. All the computations of the proof will be made assuming, first, $g \in H^1(0, T; L^2(-1, 1)) \cap L^2(0, T; H^2 \cap H_0^1(-1, 1))$. Then, the conclusion of Proposition 1 will follow by a density argument.

Consider the weight function

$$\alpha(t, v) := \frac{M\beta(v)}{t(T-t)}, \quad (t, v) \in (0, T) \times \mathbb{R}, \quad (35)$$

where $\beta \in C^2([-1, 1])$ satisfies

$$\beta \geq 1 \quad \text{on } (-1, 1), \quad (36)$$

$$|\beta'| > 0 \quad \text{on } [-1, a'] \cup [b', 1], \quad (37)$$

$$\beta'(1) > 0, \quad \beta'(-1) < 0, \quad (38)$$

$$\beta'' < 0 \quad \text{on } [-1, a'] \cup [b', 1] \quad (39)$$

and $M = M(T, n, \beta) > 0$ will be chosen later on. We also introduce the function

$$z(t, v) := g(t, v)e^{-\alpha(t, v)}, \quad (40)$$

that satisfies

$$e^{-\alpha} \mathcal{P}_n g = P_1 z + P_2 z + P_3 z, \quad (41)$$

where

$$P_1 z := -\frac{\partial^2 z}{\partial v^2} + (\alpha_t - \alpha_v^2)z, \quad P_2 z := \frac{\partial z}{\partial t} - 2\alpha_v \frac{\partial z}{\partial v} + inv^\nu z, \quad P_3 z := -\alpha_{vv}z. \quad (42)$$

We develop the classical proof (see [25]), taking the $L^2(Q)$ -norm in the identity (41), then developing the double product which leads to

$$\int_Q \left(\Re[P_1 z \overline{P_2 z}] - \frac{1}{2} |P_3 z|^2 \right) dv dt \leq \int_Q |e^{-\alpha} \mathcal{P}_n g|^2 dv dt, \quad (43)$$

where $Q := (0, T) \times (-1, 1)$. After computations (see [5] for details, we get

$$\int_Q |z|^2 \left\{ -\frac{1}{2}(\alpha_t - \alpha_v^2)_t + [(\alpha_t - \alpha_v^2)\alpha_v]_v - \frac{1}{2}\alpha_{vv}^2 \right\} dv dt + \int_Q \left\{ n\gamma v^{\gamma-1} \Im \left(\bar{z} \frac{\partial z}{\partial v} \right) - \alpha_{vv} \left| \frac{\partial z}{\partial v} \right|^2 \alpha_{vv} \right\} dv dt \leq \int_Q |e^{-\alpha} \mathcal{P}_n g|^2 dv dt. \tag{44}$$

Now, in the left hand side of (44), we separate the terms on $(0, T) \times (a', b')$ and those on $(0, T) \times [(-1, a') \cup (b', 1)]$. One has

$$-\alpha_{vv}(t, v) \geq \frac{C_1 M}{t(T-t)} \quad \forall v \in [-1, a'] \cup [b', 1], \quad t \in (0, T),$$

$$|\alpha_{vv}(t, v)| \leq \frac{C_2 M}{t(T-t)} \quad \forall v \in [a', b'], \quad t \in (0, T), \tag{45}$$

where $C_1 = C_1(\beta) := \min\{-\beta''(x); x \in [-1, a'] \cup [b', 1]\}$ is positive thanks to the assumption (39) and $C_2 = C_2(\beta) := \sup\{|\beta''(x)|; x \in [a', b']\}$. Moreover,

$$-\frac{1}{2}(\alpha_t - \alpha_v^2)_t + [(\alpha_t - \alpha_v^2)\alpha_v]_v - \frac{1}{2}\alpha_{vv}^2 = \frac{1}{(t(T-t))^3} \left\{ M\beta(3Tt - T^2 - 3t^2) + M^2 \left[(2t - T)(\beta''\beta + 2\beta'^2) - \frac{t(T-t)\beta''^2}{2} \right] - 3M^3\beta''\beta'^2 \right\}.$$

Hence, owing to (37) and (39), there exist $m_1 = m_1(\beta) > 0$ $C_3 = C_3(\beta) > 0$ and $C_4 = C_4(\beta) > 0$ such that for every $M \geq M_1$ and $t \in (0, T)$,

$$-\frac{1}{2}(\alpha_t - \alpha_v^2)_t + [(\alpha_t - \alpha_v^2)\alpha_v]_v - \frac{1}{2}\alpha_{vv}^2 \geq \frac{C_3 M^3}{[t(T-t)]^3} \quad \forall v \in [0, a'] \cup [b', 1],$$

$$\left| -\frac{1}{2}(\alpha_t - \alpha_v^2)_t + [(\alpha_t - \alpha_v^2)\alpha_v]_v - \frac{1}{2}\alpha_{vv}^2 \right| \leq \frac{C_4 M^3}{[t(T-t)]^3} \quad \forall v \in [a', b'] \tag{46}$$

where

$$M_1 = M_1(T, \beta) := m_1(\beta)(T + T^2). \tag{47}$$

Using (44), (45) and (46), we deduce, for every $M \geq M_1$,

$$\begin{aligned}
 & \int_0^T \int_{(-1,a') \cup (b',1)} \frac{C_1 M}{t(T-t)} \left| \frac{\partial z}{\partial v} \right|^2 dv dt \\
 & + \int_0^T \int_{(-1,a') \cup (b',1)} \left[\frac{C_3 M^3}{(t(T-t))^3} |z|^2 - |n|\gamma|v|^{\gamma-1} \Im \left(\frac{\partial z}{\partial v} \bar{z} \right) \right] dv dt \\
 & \leq \int_0^T \int_{a'}^{b'} \left[\frac{C_2 M}{t(T-t)} \left| \frac{\partial z}{\partial x} \right|^2 + \frac{C_4 M^3}{(t(T-t))^3} |z|^2 + |n|\gamma|v|^{\gamma-1} \Im \left(\frac{\partial z}{\partial v} \bar{z} \right) \right] dv dt \\
 & + \int_Q e^{-\alpha \mathcal{P}_n g} |z|^2 dv dt. \tag{48}
 \end{aligned}$$

Let

$$M_2 = M_2(T, \beta) := \frac{T^2 \sqrt{|n|\gamma}}{4\sqrt[4]{C_1 C_3}}. \tag{49}$$

When $M \geq M_2$, we have

$$\begin{aligned}
 \left| |n|\gamma|v|^{\gamma-1} \Im \left(\frac{\partial z}{\partial v} \bar{z} \right) \right| & \leq \frac{1}{2} \frac{C_3 M^3}{(t(T-t))^3} |z|^2 + \frac{1}{2} \frac{(t(T-t))^3}{C_3 M^3} \gamma^2 n^2 \left| \frac{\partial z}{\partial v} \right|^2 \\
 & \leq \frac{1}{2} \frac{C_3 M^3}{(t(T-t))^3} |z|^2 + \frac{C_1 M}{2t(T-t)} \left| \frac{\partial z}{\partial v} \right|^2,
 \end{aligned} \tag{50}$$

because

$$\begin{aligned}
 \frac{1}{2} \frac{(t(T-t))^3}{C_3 M^3} \gamma^2 n^2 & = \frac{C_1 M}{2t(T-t)} \frac{(t(T-t))^4 \gamma^2 n^2}{C_1 C_3 M^4} \\
 & \leq \frac{C_1 M}{2t(T-t)} \frac{(T^2/4)^4 \gamma^2 n^2}{C_1 C_3 M^4} \\
 & = \frac{C_1 M}{2t(T-t)} \frac{M_2^4}{M^4}
 \end{aligned}$$

From now on, we take

$$M = M(T, n, \beta) := C_2 \max\{T + T^2; \sqrt{|n|T^2}\} \tag{51}$$

where

$$C_2 = C_2(\beta) := \max \left\{ m_1; \frac{1}{4\sqrt[4]{C_1 C_3}} \right\}$$

so that $M \geq M_1$ and M_2 (see (47) and (49)). We have

$$\begin{aligned} & \int_0^T \int_{(0-1, a') \cup (b', 1)} \left(\frac{C_1 M}{2t(T-t)} \left| \frac{\partial z}{\partial v} \right|^2 + \frac{C_3 M^3}{2(t(T-t))^3} |z|^2 \right) dv dt \\ & \leq \int_0^T \int_{a'}^{b'} \left(\frac{C'_2 M}{t(T-t)} \left| \frac{\partial z}{\partial v} \right|^2 + \frac{C_6 M^3}{(t(T-t))^3} |z|^2 \right) dv dt \\ & \quad + \int_Q |e^{-\alpha} \mathcal{P}_n g|^2 dv dt, \end{aligned} \tag{52}$$

where $C_6 = C_6(\beta) := C_4 + C_3/2$, $C'_2 = C_2(\beta) = C_2 + C_1/2$. Since for every $\epsilon > 0$

$$\begin{aligned} & \frac{C_1 M}{2t(T-t)} \left| \frac{\partial g}{\partial x} - \alpha_x g \right|^2 + \frac{C_3 M^3}{2(t(T-t))^3} |g|^2 \\ & \geq \left(1 - \frac{1}{1+\epsilon} \right) \frac{C_1 M}{2t(T-t)} \left| \frac{\partial g}{\partial x} \right|^2 + \frac{M^3}{2(t(T-t))^3} (C_3 - \epsilon C_1(\beta')^2) |g|^2 \end{aligned} \tag{53}$$

and choosing

$$\epsilon = \epsilon(\beta) := \frac{C_3}{2C_1 \|\beta'\|_\infty^2},$$

from (52), (53) and (40) we deduce that

$$\begin{aligned} & \int_0^T \int_{(0, a') \cup (b', 1)} \left(\frac{C_7 M}{t(T-t)} \left| \frac{\partial g}{\partial v} \right|^2 + \frac{C_3 M^3 |g|^2}{4(t(T-t))^3} \right) e^{-2\alpha} dv dt \\ & \leq \int_Q |e^{-\alpha} \mathcal{P}_n g|^2 dv dt + \int_0^T \int_{a'}^{b'} \left(\frac{C_9 M^3 |g|^2}{(t(T-t))^3} + \frac{C_8 M}{t(T-t)} \left| \frac{\partial g}{\partial v} \right|^2 \right) e^{-2\alpha} dv dt, \end{aligned} \tag{54}$$

where $C_7 = C_7(\beta) := [1 - 1/(1+\epsilon)]C_1/2$, $C_8 = C_8(\beta) := 2C'_2$ and $C_9 = C_9(\beta) := C_6 + 2C'_2 \sup\{\beta'(x)^2; x \in [a', b']\}$. Adding the same quantity in both sides, we get

$$\begin{aligned} & \int_Q \left(\frac{C_7 M}{t(T-t)} \left| \frac{\partial g}{\partial v} \right|^2 + \frac{C_3 M^3 |g|^2}{4(t(T-t))^3} \right) e^{-2\alpha} dv dt \leq \int_Q |e^{-\alpha} \mathcal{P}_n g|^2 dv dt \\ & \quad + \int_0^T \int_{a'}^{b'} \left(\frac{C_{11} M^3 |g|^2}{(t(T-t))^3} + \frac{C_{10} M}{t(T-t)} \left| \frac{\partial g}{\partial v} \right|^2 \right) e^{-2\alpha} dv dt, \end{aligned} \tag{55}$$

where $C_{10} = C_{10}(\beta) := C_8 + C_7$ and $C_{11} = C_{11}(\beta) := C_9 + C_3/4$. Thanks to a cut-off function ρ such that

$$0 \leq \rho \leq 1, \quad \rho \equiv 1 \text{ on } (a', b'), \quad \text{Supp}(\rho) \subset (a, b)$$

it is classic to get

$$\begin{aligned} & \int_0^T \int_{a'}^{b'} \frac{C_{10}M}{t(T-t)} \left| \frac{\partial g}{\partial v} \right|^2 e^{-2\alpha} \, dv dt \\ & \leq \int_Q |\mathcal{P}_n g|^2 e^{-2\alpha} \, dv dt + \int_0^T \int_a^b \frac{C_{12}M^3 |g|^2 e^{-2\alpha}}{(t(T-t))^3} \, dv dt \end{aligned}$$

for some constant $C_{12} = C_{12}(\beta) > 0$. Combining (55) with the previous inequality, we get

$$\begin{aligned} & \int_Q \left(\frac{C_7M}{t(T-t)} \left| \frac{\partial g}{\partial v} \right|^2 + \frac{C_3M^3 |g|^2}{4(t(T-t))^3} \right) e^{-2\alpha} \, dv dt \\ & \leq \int_Q 2|e^{-\alpha} \mathcal{P}_n g|^2 \, dv dt + \int_0^T \int_a^b \frac{C_{13}M^3 |g|^2}{(t(T-t))^3} e^{-2\alpha} \, dv dt, \end{aligned} \tag{56}$$

where $C_{13} = C_{13}(\beta, \rho) := C_{11} + C_{12}$. Then, the global Carleman estimates (10) holds with

$$C_1 = C_1(\beta) := \frac{\min \{C_7; C_3/4\}}{\max \{2; C_{13}\}}.$$

References

1. Alabau-Boussouira F, Cannarsa P, Fragnelli G (2006) Carleman estimates for degenerate parabolic operators with applications to null controllability. *J Evol Equ* 6(2):161–204
2. Alinhac S, Zuily C (1981) Uniqueness and nonuniqueness of the cauchy problem for hyperbolic operators with double characteristics. *Commun Partial Differ Equ* 6(7):799–828
3. Almog Y (2008) The stability of the normal state of superconductors in the presence of electric currents. *SIAM J Math Anal* 40(2):824–850
4. Beauchard K, Cannarsa P, Guglielmi R (2012) Null controllability of Grushin-type operators in dimension two. *J Eur Math Soc* (to appear)
5. Beauchard K, Zuazua E (2009) Some controllability results for the 2D Kolmogorov equation. *Ann Inst H Poincaré Anal Non Linéaire* 26:1793–1815
6. Benabdallah A, Dermenjian Y, Le Rousseau J (2007) Carleman estimates for the one-dimensional heat equation with a discontinuous coefficient and applications to controllability and an inverse problem. *J Math Anal Appl* 336(2):865–887
7. Benabdallah A, Dermenjian Y, Le Rousseau J (2007) On the controllability of linear parabolic equations with an arbitrary control location for stratified media. *C R Math Acad Sci Paris* 344(6):357–362
8. Br Tzis H (1983) *Analyse fonctionnelle, theorie et applications*. Masson, Paris

9. Cannarsa P, de Teresa L (2009) Controllability of 1-d coupled degenerate parabolic equations. *Electron J Differ Equ Paper No.* 73:21
10. Cannarsa P, Fragnelli G, Rocchetti D (2007) Null controllability of degenerate parabolic operators with drift. *Netw Heterog Media* 2(4):695–715 (electronic)
11. Cannarsa P, Fragnelli G, Rocchetti D (2008) Controllability results for a class of one-dimensional degenerate parabolic problems in nondivergence form. *J Evol Equ* 8:583–616
12. Cannarsa P, Martinez P, Vancostenoble J (2004) Persistent regional null controllability for a class of degenerate parabolic equations. *Commun Pure Appl Anal* 3(4):607–635
13. Cannarsa P, Martinez P, Vancostenoble J (2005) Null controllability of degenerate heat equations. *Adv Differ Equ* 10(2):153–190
14. Cannarsa P, Martinez P, Vancostenoble J (2008) Carleman estimates for a class of degenerate parabolic operators. *SIAM J Control Optim* 47(1):1–19
15. Cannarsa P, Martinez P, Vancostenoble J (2009) Carleman estimates and null controllability for boundary-degenerate parabolic operators. *C R Math Acad Sci Paris* 347(3–4):147–152
16. Doubova A, Fernández-Cara E, Zuazua E (2002) On the controllability of parabolic systems with a nonlinear term involving the state and the gradient. *SIAM J Control Optim* 42(3):798–819
17. Doubova A, Osses A, Puel J-P (2002) Exact controllability to trajectories for semilinear heat equations with discontinuous diffusion coefficients. *ESAIM Control Optim Calc Var* 8:621–661
18. Ervedoza S (2008) Control and stabilization properties for a singular heat equation with an inverse-square potential. *Commun Partial Differ Equ* 33(10–12):1996–2019
19. Fabre C, Puel JP, Zuazua E (1995) Approximate controllability of the semilinear heat equation. *Proc Roy Soc Edinb* 125A:31–61
20. Fattorini HO, Russel D (1971) Exact controllability theorems for linear parabolic equations in one space dimension. *Arch Ration Mech Anal* 43:272–292
21. Fernández-Cara E, Zuazua E (2000) Null and approximate controllability for weakly blowing up semilinear heat equations. *Ann Inst H Poincaré Anal Non Linéaire* 17:583–616
22. Fernández-Cara E, Zuazua E (2000) The cost of approximate controllability for heat equations: the linear case. *Adv Differ Equ* 5(4–6):465–514
23. Fernández-Cara E, Zuazua E (2002) On the null controllability of the one-dimensional heat equation with BV coefficients. *Comput Appl Math* 12:167–190
24. Flores C, de Teresa L (2010) Carleman estimates for degenerate parabolic equations with first order terms and applications. *C R Math Acad Sci Paris* 348(7–8):391–396
25. Fursikov AV, Imanuvilov OY (1994) Controllability of evolution equations. *Lecture notes series, Seoul National University Research Institute of Mathematics Global Analysis Research Center, Seoul*, 34
26. González-Burgos M, de Teresa L (2007) Some results on controllability for linear and nonlinear heat equations in unbounded domains. *Adv Diff Equ* 12(11):1201–1240
27. Hörmander L (1967) Hypoelliptic second order differential equations. *Acta Math* 119:147–171
28. Imanuvilov OY (1993) Boundary controllability of parabolic equations. *Uspekhi Mat Nauk* 48(3(291)):211–212
29. Imanuvilov OY (1995) Controllability of parabolic equations. *Math Sb* 186(6):109–132
30. Imanuvilov OY, Yamamoto M (2000) Carleman estimate for a parabolic equation in sobolev spaces of negative order and its applications. In: Chen G et al (eds) *Control of nonlinear distributed parameter systems*. Marcel Dekker, New York, pp 113–137
31. Lebeau G, Robbiano L (1995) Contrôle exact de l'équation de la chaleur. *Commun Partial Differ Equ* 20:335–356
32. Lebeau G, Le Rousseau J (2012) On Carleman estimates for elliptic and parabolic operators. Applications to unique continuation and control of parabolic equations. *ESAIM:COCV* 18:712–747
33. Lopez A, Zuazua E (2002) Uniform null controllability for the one dimensional heat equation with rapidly oscillating periodic density. *Ann IHP Anal Non linéaire* 19(5):543–580
34. Martinez P, Vancostenoble J (2006) Carleman estimates for one-dimensional degenerate heat equations. *J Evol Equ* 6(2):325–362
35. Martinez P, Vancostenoble J, Raymond J-P (2003) Regional null controllability of a linearized Crocco type equation. *SIAM J Control Optim* 42(2):709–728
36. Miller L (2005) On the null-controllability of the heat equation in unbounded domains. *Bull Des Sci Math* 129(2):175–185
37. Miller L (2006) On exponential observability estimates for the heat semigroup with explicit rates. *Rendiconti Lincei: Matematica e Applicazioni* 17(4):351–366

38. Pazy A (1983) Semigroups of linear operators and applications to partial differential equations. Applied Mathematical Sciences, vol 44. Springer, New York
39. Le Rousseau J (2007) Carleman estimates and controllability results for the one-dimensional heat equation with bv coefficients. *J Differ Equ* 233(2):417–447
40. Vancostenoble J, Zuazua E (2008) Null controllability for the heat equation with singular inverse-square potentials. *J Funct Anal* 254(7):1864–1902
41. Villani C (2009) Hypocoercivity. *Mem Am Math Soc* 202(950)
42. Zuazua E (1997) Approximate controllability of the semilinear heat equation: boundary control. In: Bristeau MO et al (eds) International conference in honour of Prof. R. Glowinski, computational sciences for the 21st century. Wiley, pp 738–747
43. Zuazua E (1997) Finite dimensional null-controllability of the semilinear heat equation. *J Math Pures et Appl* 76:237–264